

## Invasive Aspergillosis in Patients with AIDS

Saye H. Khoo and David W. Denning

From the Department of Infectious Diseases and Tropical Medicine, Monsall Unit, North Manchester General Hospital, Manchester; and the Department of Medicine, University of Manchester, Hope Hospital, Salford, United Kingdom

The prolonged survival of profoundly immunocompromised patients with AIDS has contributed to the increasing recognition of aspergillus infections as an emerging problem. Nevertheless, many of these infections continue to be diagnosed only at autopsy. In this article we review details of 293 reported cases. Invasive aspergillosis occurs in advanced AIDS and most commonly affects the lungs, although brain involvement has also been frequently reported. The diagnosis is often difficult to make while the patient is alive, although examination of specimens obtained via bronchoalveolar lavage, percutaneous needle aspiration, or biopsy is often successful. Biopsy of the affected organ along with histologic examination and culture may be necessary for diagnosis. The dismal prognosis of invasive aspergillosis in patients with AIDS can be improved only with earlier diagnosis of disease and the availability of more-effective antifungal regimens.

Invasive aspergillosis, although typically occurring in the immunocompromised patient, has not been well recognized as an AIDS-associated condition. Yet the finding of invasive aspergillosis, even in early autopsy series of patients with AIDS, has been well documented [1–12], and the recent surge of case reports has highlighted infections due to *Aspergillus* species in these patients. This article reviews the epidemiology, clinical spectrum, diagnosis, and treatment of aspergillus infections in 293 patients positive for human immunodeficiency virus (HIV) whose cases have been described in the world literature [2–76]. (Whenever we have suspected duplication, we have attempted to exclude such cases.)

### Epidemiology

*Aspergillus* species have a worldwide distribution, and infections in patients with AIDS have been reported from North America, Europe, Africa, and Asia. In many developing countries, cases appear to occur less frequently among these patients, either because of underdiagnosis or as a result of reduced survival. The true incidence of aspergillus infections among HIV-positive patients is difficult to estimate. Owing to the difficulty of diagnosis of clinically unsuspected disease, the diagnosis is frequently not made until autopsy. Retrospective surveys of clinically diagnosed disease have indicated an incidence of 0.9%–8.6% [14, 27, 43, 76] among

patients with AIDS. A survey of 12 major autopsy series [1–12] demonstrated a prevalence of 0–12% (with an overall prevalence of 4%), although in five of these series, only neuropathological findings were reported [1–4, 9]. The diagnosis of aspergillus infection was made while the patient was alive in 169 (71%) of 239 clinical cases reported [8–13, 15–43, 46–62, 71, 73, 75]. These figures probably reflect a bias in favor of publishing reports of cases diagnosed while the patient was still alive. In our survey of five autopsy series involving 912 subjects, only 9 (22%) of the 41 reported cases of invasive aspergillosis were diagnosed antemortem [8–12]. It is thus probable that most cases remain undiagnosed (table 1).

Invasive disease appears to occur almost exclusively in patients with late- or end-stage HIV disease, occurring at a median of 10–26 months after the onset of AIDS [13, 15, 17, 30, 76]. Of the 293 reported cases, only four were classified as CDC Group II at presentation [13–14, 26, 39]. CD4 cell counts, when reported, have almost always been low ( $<100/\text{mm}^3$ ) [13–22, 71, 76], although disease has occasionally occurred in adults with counts of  $\geq 200/\text{mm}^3$  [37, 39, 69, 73–74]. Thus it is not surprising that intercurrent opportunistic pathogens are usually present, most frequently *Pneumocystis carinii* or other bacterial agents of pneumonia, as well as cytomegalovirus (CMV), *Mycobacterium avium* complex, or *Toxoplasma gondii*.

As with many other infections in advanced HIV disease, numerous risk factors for aspergillosis are postulated, but few are proven since data mostly derive from case reports. No case-control studies that assess relative risk have been published. Recognized risk factors for invasive aspergillosis in non-HIV-infected immunocompromised patients are neutropenia and prior corticosteroid or cytotoxic chemotherapy for malignant disease; the significance of administration of recently available broad-spectrum antibiotic therapy is un-

---

Reprints or correspondence: Dr. D. W. Denning, Department of Infectious Diseases and Tropical Medicine, University of Manchester, North Manchester Hospital, Delauney's Road, Manchester M8 6RL, United Kingdom.

**Clinical Infectious Diseases** 1994;19(Suppl 1):S41–8  
 © 1994 by The University of Chicago. All rights reserved.  
 1058–4838/94/1902–0006\$02.00

**Table 1.** Prevalence of invasive aspergillosis at autopsy among 1,569 patients.

Reference	No. of autopsies performed	No. of cases of aspergillosis detected	Frequency (%)
[1]	29	0*	0
[2]	153	7*	5
[3]	221	2*	1
[4]	135	1*	1
[5]	56	5	9
[6]	13	1	7
[7]	50	3	6
[8]	12	1	8
[9]	40	1*	3
[10]	101	7	7
[11]	529	26	12
[12]	230	5	2
Total number of cases	1,569	59	4

\* Only neuropathological examination results reported.

certain. Neutropenia ( $<1,000$  neutrophils  $\times 10^6/L$ ) was present in 92 (46%) of 202 evaluable patients with invasive aspergillosis [7, 11–44, 71, 76]; a neutrophil count of  $<500 \times 10^6/L$  was documented in 6%–50% of patients in some series [13, 15, 17]. Patients may present with invasive aspergillosis even after their neutrophil counts have returned to normal (because of recovery of bone marrow from the effects of cytotoxic agents or zidovudine). Corticosteroid therapy was documented in 79 (39%) of the 202 cases; doses, where specified, were  $\geq 10$  mg of prednisolone/d or its equivalent [12, 15–16, 40–42]. Prior broad-spectrum antibiotic therapy was documented in 64 (32%) of the cases; this figure is probably underestimated. The significance of other factors is less clear. The speculation that inhalation of marijuana [13, 14] contaminated with *Aspergillus* species might constitute a risk, though intriguing, remains unproven. There are also possible links with alcohol consumption [20, 26, 27].

Among patients with invasive pulmonary aspergillosis, the apparently high frequency of preceding or intercurrent pneumonia due to *P. carinii* and other bacteria (present in 124 [73%] of 169 reported cases [7, 11–20, 22–23, 25–33, 35–36, 40, 44, 71, 75]) suggests that these infections may also be a contributory factor. However, conclusive evidence is lacking since pneumonia is a common event in patients with advanced-stage HIV disease. The development of aspergillomas in some HIV-infected patients, as is true for other groups of patients such as those with pulmonary sarcoidosis, is facilitated by existing lung damage [61, 68, 76] and has occurred within a lung cyst following pneumocystis pneumonia [57].

Some patients have no apparent risk factors for aspergillosis apart from HIV infection itself [17, 72]. Contrary to the conclusions drawn in earlier reports [7, 11], it is likely that

advanced HIV infection constitutes an independent risk factor for the development of invasive aspergillosis. Alveolar macrophage function constitutes the first line of defense against *Aspergillus* conidia, and neutrophils are the major component (with some contribution from monocytes and complement activation) defending against *Aspergillus* hyphae; the function and numbers of these cells may be affected by advanced HIV infection [77]. In studies of HIV-infected children, the activity of neutrophils and peripheral blood macrophages against *Aspergillus fumigatus* was impaired [78–79], suggesting that there is increased susceptibility to infection with this organism, even if absolute numbers of cells are maintained.

Aspergilli are common environmental saprophytes, accounting for up to 40% of the fungal flora in hospital and home environments. Many nosocomial outbreaks in other groups have been described (e.g., bone marrow transplant recipients). A cluster of seven cases occurring on an AIDS ward has been described [36]; nosocomial spread was suspected but remains unproven.

In some of the reports surveyed, no details were available on the species of *Aspergillus* isolated. In others, the diagnosis was made histologically. While it may be difficult to differentiate *Aspergillus* hyphae from those of other pathogenic molds, we consider the overwhelming majority of these cases do indeed represent aspergillosis. The most common species causing disease in patients with AIDS is *A. fumigatus*, which represented 73 (84%) of 87 isolates recovered [12, 13, 15–17, 20, 23–24, 26, 29–30, 33–34, 36–37, 39–40, 42, 48–49, 52, 54, 57, 68–71, 73–75]. *Aspergillus flavus* was identified in 7 cases (8%), *Aspergillus niger* in 4 (5%), and *Aspergillus terreus* in 3 (3%) [12, 15–17, 40, 53, 62, 69, 75].

### Clinical Syndromes

Table 2 lists the major sites of infection. The lung is the organ most frequently involved ( $>70\%$  of cases); brain involvement was reported in 10% of cases. Disease is almost always invasive, although aspergillus bronchitis and aspergillomas have been reported. Whether or not there is increased risk of allergic disease, such as allergic bronchopulmonary aspergillosis or allergic aspergillus sinusitis, is unknown.

The diagnosis of infection while the patient is alive is often possible only with culture and/or histologic examination of a biopsy specimen from the affected organ. Results of blood cultures, although only rarely positive [80], should suggest the possibility of disseminated disease [20], fungal endocarditis or myocarditis [37], mycotic aneurysm, or an infected central venous catheter. Precipitins (*Aspergillus*-specific IgG) are elevated only in patients with aspergilloma [61]. In other groups of immunosuppressed patients, *Aspergillus* antigen in blood or urine may be detected occasionally [81], but in general, serology is unhelpful.

**Table 2.** Sites of aspergillosis in 293 cases among HIV-positive patients.

Site/infection	No. of patients (%) with indicated infection	References
<b>Respiratory tract*</b>		
Sinusitis	9 (3)	[13, 15, 17, 24, 33, 41, 53, 62, 67]
Otomastoiditis	5 (2)	[17, 48, 56]
Laryngitis	2 (<1)	[23, 33]
Tracheobronchitis	11 (4)	[13, 17, 30, 53, 71-72]
Obstructing bronchial aspergillosis	5 (2)	[13, 32, 39]
Invasive pulmonary aspergillosis	208 (71)	[7, 9, 11-23, 25-32, 35, 36, 40, 43, 46, 50-53, 58, 59, 63, 66, 69, 70, 74-76]
Empyema/pleural mass	5 (2)	[13, 30, 36, 69]
Aspergilloma	4 (1)	[57, 61, 68, 76]
<b>CNS*</b>		
Brain†	30 (10)	[2-4, 9, 11, 13, 15, 17, 26, 30, 43-44, 51-52, 54, 60, 64, 67, 69-70, 75]
<b>Other systems*</b>		
Cardiac	10 (3)	[15, 17, 28, 29, 31, 44, 54, 67, 69]
Renal	12 (4)	[11, 15, 17, 18, 31, 34, 49, 54, 66, 69]
Thyroid	4 (1)	[11, 31]
Other*: spleen (3), musculoskeletal (3), lymph node (2), pancreas (2), liver (2), mouth, palate, adrenal, epididymis, gastrointestinal, skin	16 (5)	[8, 13, 17, 18, 20, 23, 30, 36, 38, 42, 47, 54, 55, 71, 73]
Two or more organs involved	47 (16)	[4, 10-11, 13, 15-18, 20, 23, 26, 28, 29, 32, 33, 36, 41, 44, 51-54, 69, 73]
Site of disease not specified	13 (4)	[3, 45, 64]

\* Includes disease occurring in more than one site.


† Including spinal cord involvement in one patient.

**Respiratory tract infections.** Invasive pulmonary disease presented most frequently as cough (92%), fever (91%), dyspnea (65%), chest pain (24%), and hemoptysis (9%) in our survey of 78 evaluable patients [13, 17-19, 26, 31, 32, 35, 40, 46, 69]. There is often a prior history of pneumonia due to *Pneumocystis* or other bacteria or CMV. The chest radiograph almost always shows abnormalities, but appearances may be variable. They include localized or diffuse infiltrates and nodules. Cavitory lesions with thick or thin walls are relatively distinctive. They are usually present in the upper lobes (figure 1) and may be confused with those of tuberculosis. Pleural effusions may occasionally be present; two cases of empyema due to *Aspergillus* species have been reported [36, 69].

The diagnosis of invasive pulmonary aspergillosis can be difficult. *Aspergillus* species may be present in the respiratory tract as commensals in both healthy individuals and the immunocompromised [82], including ~4% of patients with AIDS [12, 83-84]. Thus, the isolation of *Aspergillus* species from respiratory secretions does not automatically imply invasive disease. However, studies of bone marrow transplant recipients as well as of those with a hematologic malignancy have shown that positive sputum culture is strongly predictive of invasive disease [82, 85]. The presence of *Aspergillus*

species in the sputum of immunocompromised patients, including those with AIDS, in association with abnormal findings on a chest radiograph should suggest the possibility of invasive pulmonary aspergillosis [12]. The converse is not true: a negative sputum culture does not exclude the possibility of aspergillosis. Bronchoscopic findings may appear to be normal, but lavage usually yields the organism [17]. Transbronchial biopsy (although often unrewarding) and transthoracic needle aspiration of localized lesions may also clinch the diagnosis [13].

The three patients with aspergilloma who have been described all had existing lung damage (one had aspergilloma within a cyst that had resulted from pneumocystis pneumonia). Characteristic changes were seen on radiographs and computed tomographic (CT) scans of the thorax. Hemoptysis may be fatal [68] and require surgical intervention. Ulcerative or pseudomembranous tracheobronchitis caused by *Aspergillus* species is a recently described entity occurring in immunocompromised patients who do not have AIDS [86-87] and has been reported in 11 patients with AIDS [13, 17, 30, 71-72]. Patients present with dyspnea and wheezing (which may mimic asthma); fever and a cough (or hemoptysis) may also be present. Bronchoscopy reveals ulcerative lesions or fungal necrotic pseudomembranes [13, 17] that



**Figure 1.** Finding of pulmonary aspergillosis in a patient who had AIDS for 3 years (CD4 count,  $>10/\text{mm}^3$ ). Right middle-lobe consolidation and a left upper-lobe cavity as seen in a previously normal lung; the diagnosis was made by culture of a bronchoalveolar lavage specimen and was confirmed at autopsy.

may be extensive and invade peribronchial tissue and local blood vessels, resulting in distant spread of disease. Associated lung parenchymal disease is found in 50% of cases, and distant dissemination is found in 25% of cases [71].

A noninvasive form of airway disease, called obstructing bronchial aspergillosis, has also been described in patients with AIDS [13, 32, 39]; a similar case occurred in a bone marrow transplant recipient [88]. Symptoms are cough, fever, wheezing, hemoptysis, and hypoxemia. The distinctive feature is that large mucoid casts full of *Aspergillus* organisms are expectorated spontaneously or are encountered at bronchoscopy. These casts plug the airways and may result in segmental or lobar atelectasis, usually affecting the lower lobes. There is little or no bronchial inflammation in the early stages. Chest radiographs show bilateral lower-lobe or generalized hazy shadowing. Disease may become invasive and extend upwards to produce pseudomembranous tracheobronchitis, particularly if the disease is left untreated [13]. The diagnosis of aspergillus tracheobronchitis and obstructing bronchial aspergillosis is established at bronchoscopy.

The larynx was the site of disease in two patients [23, 33] who presented with hoarseness and/or aphonia. The diagno-

sis was made by direct laryngoscopy or bronchoscopy and was confirmed by histologic and microbiological examination of biopsy material. In laryngeal aspergillosis the cords become thickened and may be destroyed or covered by an extensive membrane.

Nine cases of aspergillosis of the sinuses in patients with AIDS have been reported; the patients usually presented with nasal discharge and facial pain. These were all invasive infections, sometimes penetrating into adjacent tissue such as the orbit and brain [24, 41, 67]. The sphenoid sinuses have also been affected (figure 2) Diagnosis is made by culture and/or biopsy, and in addition to specific antifungal chemotherapy, surgical debridement may be necessary to control disease.

Invasive otomastoiditis (invasive external otitis) typically presents as ear pain (often severe) and otorrhea; auroscopic examination reveals a perforated tympanic membrane [17, 48, 56]. The mastoid air cells are usually involved, but CT scans of the affected area do not always reveal abnormality. Facial nerve involvement has been documented. Culture is mandatory, since other organisms (including fungi such as *Pseudallescheria*) are also recognized causes of disease.

*CNS infections.* All too often CNS involvement is documented only at autopsy. CNS disease often takes the form of cerebral infarction, hemorrhage, or abscess formation; the patient presents with fever and seizures or symptoms mimicking those associated with stroke or a space-occupying lesion. Spinal cord involvement and chronic meningitis have been documented [69]. A CT scan of the head may show single or multiple nonenhancing lesions, often with surrounding edema; in addition there may be bony invasion [69, 75]. Examination of a CSF sample is usually unhelpful. The diagnosis is difficult to make since diverse pathology produces



**Figure 2.** CT scan showing sphenoid sinus aspergillosis with destruction of the lateral wall of the sinus and temporal lobe extension. The patient had suffered from chronic headaches for 4 months; the diagnosis was made by culture of *A. fumigatus* from a percutaneous aspirate obtained through a burr hole and was confirmed at autopsy.

similar symptoms and signs in patients with AIDS. The possibility of cerebral infection is suggested by the presence of concurrent invasive pulmonary aspergillosis. If the diagnosis is in doubt, biopsy or aspiration of a localized lesion should be considered.

**Cardiac aspergillosis.** Cardiac involvement usually takes the form of endocarditis or myocarditis, manifesting as fever, cerebral embolic disease, cardiac failure, and conduction defects such as arrhythmias and atrioventricular block. The diagnosis is seldom made while the patient is alive. Positive results of fungal blood cultures should alert the clinician [37], and echocardiography may reveal vegetations [28]. Hematogenous dissemination will often result in involvement of other organs, predominantly the lung and brain.

**Other sites.** Renal aspergillosis has been described. This may take the form of invasive disease (presenting as fever, loin pain, renal dysfunction, hematuria, and pyuria [17, 18, 49]) or fungal balls [34]; surgical intervention (nephrectomy or drainage) may be required to relieve obstruction or remove infected tissue [49, 66]. A number of other organs have been infected, including the liver [47, 73] and spleen [7, 54, 73] (complicated by chylous ascites [73]), as well as the palate and mouth [18, 55], adrenal glands [23], and thyroid [11, 31]. There have also been single case reports of infection occurring in peripancreatic tissue [38], skin [8], epididymis [36], muscle [42], and prepatellar panniculitis [30]. There are no particularly distinctive clinical features, and biopsy or aspiration of the affected organ is necessary for diagnosis.

Disseminated disease affecting two or more organs is relatively common (it is present in at least 16% of cases). Again, the lungs are most frequently involved.

### Antifungal Therapy

The response to antifungal chemotherapy of AIDS patients with invasive aspergillosis is considerably inferior to that observed in patients who do not have AIDS. This finding probably reflects the advanced stage of underlying disease and the presence of intercurrent illness due to other opportunistic pathogens or tumors. Amphotericin B with or without flucytosine has been used with varying success [13, 18, 57, 89]. Although the optimal dose has yet to be defined, we suggest starting with 0.8–1.0 mg/(kg·d) and titrating the dose according to response or evidence of toxicity. The role of liposomal preparations (e.g., AmBisome; Vestar, San Dimas, CA) is even less certain, but they may be considered in cases of drug intolerance. Higher doses can and should be given (e.g., 1–4 mg/(kg·d) for invasive aspergillosis [90]). The effectiveness of amphotericin B may be limited by its toxicity as well as the associated hazards and inconvenience of long-term intravenous infusions, since lifelong therapy is usually indicated.

Response to amphotericin B therapy may be inferior to that observed when itraconazole is given with or without

flucytosine [21, 24]. If itraconazole is used, loading doses (e.g., 600 mg/d for 4 days) should be given, followed by maintenance with at least 400 mg/d administered with food. Therapy should be continued indefinitely. However, absorption of itraconazole may be impaired, and it often interacts with other drugs (particularly inducers of the p450 enzymes, such as rifampin). Serum concentrations of itraconazole should be measured to ensure absorption and to monitor levels, particularly if there are interactions accelerating its metabolism. While response to itraconazole therapy has been observed in a few patients [21–33], in many others there is neither dramatic response nor deterioration, and 25%–35% of patients may remain stable for some weeks before therapy subsequently fails or they die of another AIDS-related problem [24]. Thus, the general outlook is dismal; death occurs at a median of 2–4 months after diagnosis [13, 14, 17, 18, 23], reflecting in part the advanced stage of AIDS. Even if control of disease is successful (sustained in a minority of patients for >12 months), recrudescence is common. Hemoptysis is also a frequent fatal event in patients who have cavitory disease [13, 76]. Given the limited success with current antifungal agents, the use of newer agents as experimental therapy is appropriate for those in whom conventional treatment has failed.

Patients with obstructing bronchial aspergillosis generally respond to therapy with itraconazole [13], which may prevent progression to pseudomembranous aspergillar tracheobronchitis. Bronchoscopy may be both diagnostic and therapeutic in these individuals, especially when there is atelectasis due to obstructing fungal plugs.

### Prophylaxis

Should antifungal prophylaxis be considered and, if so, for whom? This issue remains unresolved. The decision to embark upon prophylaxis for aspergillosis in a patient with AIDS is influenced by the likelihood of disease and the effectiveness of the agent used; data on both are lacking. Although advanced HIV infection itself constitutes a risk factor, most individuals with low CD4 cell counts will not develop aspergillosis, and the cost-benefit ratio needs to be carefully evaluated, especially as therapy may be lifelong. A better option might be to target the highest risk groups; these would include patients with low CD4 cell counts who have *Aspergillus* species isolated from sputum and may also include those with neutropenia or who are receiving corticosteroid therapy. It is also unclear which drug or dose should be recommended for this purpose. The only effective oral agent presently available is itraconazole, which also has activity against species of *Candida*, *Histoplasma*, and *Cryptococcus*. However, its use in this context has not been studied and cannot be recommended at present.

Despite recent advances in antifungal therapy, the mortality associated with invasive aspergillosis remains high, even

when the infection is treated aggressively. There is clearly an urgent need for earlier diagnosis and improved drug therapy for invasive aspergillosis in patients with AIDS.

## References

- Anders K, Steinsapir KD, Iverson DJ, et al. Neuropathologic findings in the acquired immunodeficiency syndrome (AIDS). *Clin Neuropathol* 1986;5:1-20.
- Petito CK, Cho E-S, Lemann W, Navia BA, Price RW. Neuropathology of acquired immunodeficiency syndrome (AIDS): an autopsy review. *J Neuropathol Exp Neurol* 1986;45:635-46.
- Kure K, Llana JF, Lyman WD, et al. Human immunodeficiency virus-1 infection of the nervous system: an autopsy study of 268 adult, pediatric, and fetal brains. *Hum Pathol* 1991;22:700-10.
- Lang W, Miklossy J, Deruaz JP, et al. Neuropathology of the acquired immune deficiency syndrome (AIDS): a report of 135 consecutive autopsy cases from Switzerland. *Acta Neuropathol (Berl)* 1989;77:379-90.
- Niedt GW, Schinella RA. Acquired immunodeficiency syndrome. Clinicopathologic study of 56 autopsies. *Arch Pathol Lab Med* 1985;109:727-34.
- Guarda LA, Luna MA, Smith JL, Mansell PWA, Gyorkey F, Roca AN. Acquired immune deficiency syndrome: postmortem findings. *Am J Clin Pathol* 1984;81:549-57.
- Grosse G, Staib F, Seibold M, Heise W, Artigas J. Invasive aspergillosis—an AIDS-typical opportunistic infection? [abstract no M.B.P.95]. In: Program and abstracts of the 5th International Conference on AIDS (Montreal). Ottawa: International Development Research Centre, 1989.
- Hui AN, Koss MN, Meyer PR. Necropsy findings in acquired immunodeficiency syndrome. A comparison of premortem diagnoses with postmortem findings. *Hum Pathol* 1984;15:670-76.
- Gray F, Gherardi R, Keohane C, Favolini M, Sobel A, Poirier J. Pathology of the central nervous system in 40 cases of acquired immune deficiency syndrome (AIDS). *Neuropathol Appl Neurobiol* 1988;14:365-80.
- Wilkes MS, Fortin AH, Felix JC, Godwin TA, Thompson WG. Value of necropsy in acquired immunodeficiency syndrome. *Lancet* 1988;2:85-8.
- Just-Neubling G, Ganger G, Keul HG, et al. Invasive *Aspergillus* infections (IA) in AIDS: a rising problem in Frankfurt [abstract PoB 3337]. In: Program and abstracts of the 8th International Conference on AIDS (Amsterdam). Amsterdam: Congrex Holland BV, 1992.
- Pursell KJ, Telzak EE, Armstrong D. *Aspergillus* species colonization and invasive disease in patients with AIDS. *Clin Infect Dis* 1992;14:141-8.
- Denning DW, Follansbee SE, Scolaro M, Norris S, Edelstein H, Stevens DA. Pulmonary aspergillosis in the acquired immunodeficiency syndrome. *N Engl J Med* 1991;324:654-62.
- Tumbarello M, Ventura G, Caldarola G, Morace G, Cauda R. Predisposing factors and survival of pulmonary aspergillosis in AIDS patients [abstract PO-B09-1382]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). 1993.
- Minamoto GY, Barlam TF, Vander Els NJ. Invasive aspergillosis in patients with AIDS. *Clin Infect Dis* 1992;14:66-74.
- Decker CF, Parenti DM. Invasive aspergillosis in patients with HIV infection: report of two patients and a review of the literature. *J Acquir Immune Defic Syndr* 1991;4:603-6.
- Lortholary O, Meyohas MC, Dupont B, et al. Invasive aspergillosis in patients with acquired immunodeficiency syndrome: report of 33 cases. *Am J Med* 1993;95:177-87.
- Sambeat MA, Gurgui M, Ausima V, et al. Pulmonary aspergillosis in patients with AIDS [abstract PoB 3248]. In: Program and abstracts of the VIIIth International Conference on AIDS (Amsterdam). Amsterdam: Congrex Holland BV, 1992.
- Froland SS, Oppedal B, Digranes S, et al. Acquired immunodeficiency syndrome (AIDS). Clinical, immunological, pathological, and microbiological studies of the first case diagnosed in Norway. *Scand J Gastroenterol* 1985;107:82-93.
- Knani L, Maslo C, Lucet JC, et al. Overwhelming aspergillosis with positive blood cultures in an AIDS patient [abstract PuB 7286]. In: Program and abstracts of the 8th International Conference on AIDS (Amsterdam). Amsterdam: Congrex Holland BV, 1992.
- Arcidiacono MI, Cantaluppi P. Itraconazole in pulmonary aspergillosis [abstract PO-B09-1388]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). 1993.
- Wood C, Keating J, Nelson MR, Rogers T, Hawkins D, Gazzard BG. Pulmonary aspergillosis in 4 patients with advanced AIDS [abstract no PO-B09-1363]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). 1993.
- Fairley CK, Kent SJ, Street A, Mijch A, Hoy J. Invasive aspergillosis in AIDS. *Aust N Z J Med* 1991;21:747-9.
- Denning DW, Pappas PG, Kauffman CA, et al. Oral itraconazole (IZ) therapy of invasive aspergillosis (IA) [abstract no 1158]. In: Program and abstracts of the 31st Interscience Conference on Antimicrobial Agents and Chemotherapy (Chicago). Washington, DC, 1991.
- Jones PG, Cohen RL, Batts DH, Silva J. Disseminated histoplasmosis, invasive pulmonary aspergillosis, and other opportunistic infections in a homosexual patient with acquired immunodeficiency syndrome. *Sex Transm Dis* 1983;10:202-4.
- Asnis DS, Chitkara RK, Jacobson M, Goldstein JA. Invasive aspergillosis: an unusual manifestation of AIDS. *N Y State J Med* 1988;88:653-5.
- Libanore M, Pantaleoni M, Montanari P, Ghinelli F. Elevated incidence of bronchopulmonary aspergillosis (ASP-BP) in AIDS patients in northeastern Italy [abstract no PuB 7203]. In: Program and abstracts of the 8th International Conference on AIDS (Amsterdam). Amsterdam: Congrex Holland BV, 1992.
- Henochowicz S, Mustafa M, Lawrinson WE, Pistole M, Lindsay J. Cardiac aspergillosis in acquired immune deficiency syndrome. *Am J Cardiol* 1985;55:1239-40.
- Fournier JP, Bernardin G, Hoffman P, et al. Cardiopulmonary aspergillosis in an AIDS patient [abstract P32]. Third Symposium on Topics in Mycology: Mycoses in AIDS patients (Paris). 1989.
- Marlowe S, Loveless M, Ward D, et al. Invasive aspergillosis (IA) in AIDS [abstract POB09-1417]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). 1993.
- Klapholz A, Salomon N, Perlman DC, Talavera W. Aspergillosis in the acquired immunodeficiency syndrome. *Chest* 1991;100:1614-8.
- Januszewska M, Staples C, Lawson L, Roscoe DL, Montaner JGS, Phillips P. Aspergillosis: an opportunistic infection of advanced HIV disease [abstract PoB09 1376]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). 1993.
- Zylberberg L, Havard S, Gorin I, Lacau St Guily J, Benoudiba D, Escande JP. Laryngeal aspergillosis. A case treated with itraconazole [abstract PoB09-1396]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). 1993.
- Halpern M, Szabo S, Hochberg E, et al. Renal aspergilloma: an unusual cause of infection in a patient with the acquired immunodeficiency syndrome. *Am J Med* 1992;92:437-40.
- Masur H, Michelis MA, Wormser GP, et al. Opportunistic infection in previously healthy women. Initial manifestations of a community-acquired cellular immunodeficiency. *Ann Intern Med* 1982;97:533-9.
- Wright EJ, Eisen D, Hurley J, Mijch AM, Hoy JF, Lucas CR. A cluster of seven cases of *Aspergillus fumigatus* occurring in AIDS patients [abstract PuB 7254]. In: Program and abstracts of the 8th International Conference on AIDS (Amsterdam). Amsterdam: Congrex Holland BV, 1992.

- tional Conference on AIDS (Amsterdam). Amsterdam: Congrex Holland BV, 1992.
37. Schonheyder H, Hoffmann S, Jensen HE, Hansen BF, Franzmann MB. *Aspergillus fumigatus* fungaemia and myocarditis in a patient with acquired immunodeficiency syndrome. *APMIS* 1992;100:605-8.
  38. Bhatt B, Cappell MS. A peripancreatic abscess containing *Aspergillus* in a patient with the acquired immune deficiency syndrome [letter]. *Am J Gastroenterol* 1990;85:1200.
  39. Le Gros V, Stern M. Aspergillose bronchique obstructive d'evolution favorable sous corticoïdes chez un sujet porteur du virus HIV [letter]. *Ann Med Interne Paris* 1988;139:285-6.
  40. Singh N, Yu VL, Rihs JD. Invasive aspergillosis in AIDS. *South Med J* 1991;84:822-7.
  41. Schnutgen M, Hohler T, Mayet WJ, Meyer zum Buschenfelde KH. *Aspergillus* sinusitis in AIDS with rapid invasion of the orbita and the brain [abstract PoB09-1406]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). 1993.
  42. LeBras P, Zucman D, Laurian Y, Delouis C, Bouree P. Muscular aspergillosis: an unusual infection in patients with AIDS [abstract PoB09-1402]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). 1993.
  43. Rodriguez-Arroondo F, Arrizabalaga J, Irribarren JA, Garde C, Grocin AI. Invasive *Aspergillosis*: clinical and therapeutic aspects in AIDS patients [abstract P33]. In: Program and abstracts of the 3rd European Conference on Clinical Aspects and Treatment of HIV Infection (Paris). 1992.
  44. Laulund S, Visfeldt J, Klinken L. Patho-anatomical studies in patients dying of AIDS. *Acta Pathol Microbiol Immunol Scand [A]* 1986;94:201-21.
  45. Rolston KV, Radentz S, Rodriguez S. Bacterial and fungal infections in patients with the acquired immunodeficiency syndrome. *Cancer Detect Prev* 1990;14:377-81.
  46. Lissen E, Wichmann I, Jimenez JM, Andreu-Kern F. AIDS in haemophilia patients in Spain [letter]. *Lancet* 1983;1:992-3.
  47. Filice C, Brunetti E, Carnevale G, Dughetti S, Pirola F, Rondanelli EG. Ultrasonographic and microbiological diagnosis of mycotic liver abscesses in patients with AIDS. *Microbiologica* 1989;12:101-4.
  48. Reiss P, Hadderingh R, Schot LJ, Danner SA. Invasive external otitis caused by *Aspergillus fumigatus* in two patients with the acquired immunodeficiency syndrome [abstract P5]. In: Program and abstracts. Trends in the Management of Systemic Fungal Infections (Nijmegen, the Netherlands). Nijmegen, the Netherlands: Imedex, 1991.
  49. Weiss L, Picketty C, George F, Lavarde V, Kazatchkine MD. An unusual isolated kidney localisation of invasive aspergillosis in an AIDS patient [abstract PuB 7585]. In: Program and abstracts of the 8th International Conference on AIDS (Amsterdam). Amsterdam: Congrex Holland BV, 1992.
  50. Bonatz K, Weiss A, Hehlmann R, Assmus HP, Heine M. Gram-negative bacterial pneumonia with secondary aspergillosis in an AIDS patient. *Klin Wochenschr* 1991;69:853-6.
  51. Britton CB, Miller JR. Neurologic complications in acquired immunodeficiency syndrome (AIDS). *Neurol Clin* 1984;2:315-339.
  52. Hofflin JM, Remington JS. Tissue culture isolation of *Toxoplasma* from blood of a patient with AIDS. *Ann Intern Med* 1985;145:925-6.
  53. Pervez NK, Kleinerman J, Kattan M, et al. Pseudomembraneous necrotizing bronchial aspergillosis: a variant of invasive aspergillosis in a patient with hemophilia and acquired immune deficiency syndrome. *Am Rev Respir Dis* 1985;131:961-3.
  54. Cox JN, di-Dio F, Pizzolato GP, Lerch R, Pochon N. *Aspergillus* endocarditis and myocarditis in a patient with the acquired immunodeficiency syndrome (AIDS). A review of the literature. *Virchows Arch A Pathol Anat Histopathol* 1990;417:255-9.
  55. Rubin MM, Jui V, Sadoff RS. Oral aspergillosis in a patient with acquired immunodeficiency syndrome. *J Oral Maxillofac Surg* 1990;48:997-9.
  56. Strauss M, Fine E. *Aspergillus* otomastoiditis in acquired immunodeficiency syndrome. *Am J Otol* 1991;12:49-53.
  57. Torrents C, Alvarez-Castells A, de Vera PV, Coll S, Solduga C, Puy R. Postpneumocystis aspergilloma in AIDS: CT features. *J Comput Assist Tomogr* 1991;15:304-7.
  58. McLeod DT, Neill P, Robertson VJ, et al. Pulmonary diseases in patients infected with the human immunodeficiency virus in Zimbabwe, Central Africa. *Trans R Soc Trop Med Hyg* 1989;83:694-7.
  59. Marchevsky A, Rosen MJ, Chrystal G, Kleinerman J. Pulmonary complications of the acquired immunodeficiency syndrome: a clinicopathologic study of 70 cases. *Hum Pathol* 1985;16:659-70.
  60. Smith E, Pers C, Mathiesen L. Cerebral toxoplasmosis in Danish AIDS patients [abstract no. W.B.P. 27]. In: Program and abstracts of the 5th International Conference on AIDS (Montreal). Ottawa: International Development Research Centre, 1989.
  61. Hohler T, Schnutgen M, Mayet WJ, Meyer zum Buschenfelde KH. Pulmonary aspergilloma in AIDS [abstract PoB09-1410]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). 1993.
  62. Frazier R, Gathe J Jr, Stool E, LeBlanc M, Nichols M, Flaitz C. Head and neck zygomycete/*Aspergillus* infections in patients with AIDS [abstract PoB09-1365]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). 1993.
  63. Abe T. Clinical, immunological, and virological aspects in Japanese hemophiliacs and AIDS patients. *AIDS Res* 1986;2(suppl 1):S141-6.
  64. Berger JR, Moskowitz L, Fischl M, Kelley RE. The neurologic complications of AIDS: frequently the initial manifestation. *Neurology* 1984;34(suppl 1):134-35.
  65. Morrison DL, Granton JT, Kesten S, Balter MS. Cavitory aspergillosis as a complication of AIDS. *Can Assoc Radiol J* 1993;44:35-8.
  66. Valle-Gerhold J, Monzon-Alebesque F, Lopez-Lopez JA. Tratamiento percutaneo de absceso renal micotico por *Aspergillus fumigatus* en un paciente con SIDA. *Actas Urol Esp* 1992;16:492-4.
  67. Carrazana EJ, Rossitch E Jr, Morris J. Isolated central nervous system aspergillosis in the acquired immunodeficiency syndrome. *Clin Neurol Neurosurg* 1991;93:227-30.
  68. Lombardo GT, Anandarao N, Ching-Shen L, Abbate A, Becker WH. Fatal hemoptysis in a patient with AIDS-related complex and pulmonary aspergilloma. *NY State J Med* 1987;87:306-8.
  69. Woods GL, Goldsmith JC. *Aspergillus* infection of the central nervous system in patients with acquired immunodeficiency syndrome. *Arch Neurol* 1990;47:181-4.
  70. Gapen P. Neurological complications now characterizing many AIDS victims [news]. *JAMA* 1982;248:2941-2.
  71. Kemper CA, Hostetler JS, Follansbee SE, et al. Ulcerative and plaque-like tracheobronchitis due to infection with *Aspergillus* in patients with AIDS. *Clin Infect Dis* 1993;17:344-52.
  72. Salomon N, Perlman DC, Saltzman BR, Klapholz A, Talavera W. Invasive aspergillosis in AIDS: clinical manifestations and predisposing factors [abstract no M.B.P. 90]. In: Program and abstracts of the 5th International Conference on AIDS (Montreal). Ottawa: International Development Research Centre, 1989.
  73. Soriano V, Cobo J, Gutierrez M, Castejon A, Laguna F, Gonzalez-Lahoz J. *Aspergillus* hepatosplenic abscess and chylothous ascites: successful treatment with fluconazole. *European Journal of Medicine* 1992;1:253-4.
  74. Wright JL, Lawson L, Chan N, Filipenko D. An unusual form of pulmonary aspergillosis in two patients with the acquired immunodeficiency syndrome. *Am J Clin Pathol* 1993;100:57-9.
  75. Wright M, Fikrig S, Haller JO. Aspergillosis in children with acquired immune deficiency. *Pediatr Radiol* 1993;23:492-4.

76. Miller WT Jr, Sais GJ, Frank I, Geftter WB, Aronchick JM, Miller WT. Pulmonary aspergillosis in patients with AIDS: clinical and radiographic correlations. *Chest* **1994**;105:37-44.
77. Pitrak DL, Bak PM, DeMarais P, Novak RM, Andersen BR. Depressed neutrophil superoxide production in human immunodeficiency virus infection. *J Infect Dis* **1993**;167:1406-10.
78. Roilides E, Holmes A, Blake C, Pizzo PA, Walsh TJ. Impairment of neutrophil antifungal activity against hyphae of *Aspergillus fumigatus* in children infected with human immunodeficiency virus. *J Infect Dis* **1993**;167:905-11.
79. Roilides E, Holmes A, Blake C, Pizzo PA, Walsh TJ. Defective antifungal activity of monocyte-derived macrophages from human immunodeficiency virus-infected children against *Aspergillus fumigatus*. *J Infect Dis* **1993**;168:1562-5.
80. Martino P, Girmenia C, Micozzi A, et al. Fungemia in patients with leukemia. *Am J Med Sci* **1993**;306:225-32.
81. Haynes KA, Latge JP, Rogers TR. Detection of *Aspergillus* antigens associated with invasive infection. *J Clin Microbiol* **1990**;28:2040-4.
82. Barnes AJ, Denning DW. *Aspergilli*—significance as pathogens. *Rev Med Microbiol* **1993**;4:176-80.
83. Staib F, Seibold M, Grosse G. *Aspergillus* findings in AIDS patients suffering from cryptococcosis. *Mycoses* **1989**;32:516-23.
84. Daleine G, Salmon D, Lucet JF, et al. Frequence de l'isolement broncho-pulmonaire d'*Aspergillus* species chez des patients infecté par le virus de l'immunodeficiency humaine: role pathogene? *Pathol Biol Paris* **1993**;41:237-41.
85. Yu VL, Muder RR, Poorsattar A. Significance of isolation of *Aspergillus* from the respiratory tract in the diagnosis of invasive pulmonary aspergillosis. Results from a three-year prospective study. *Am J Med* **1986**;81:249-54.
86. Kramer MR, Denning DW, Marshall SE, et al. Ulcerative tracheobronchitis after lung transplantation. A new form of invasive aspergillosis. *Am Rev Respir Dis* **1991**;144:552-6.
87. Young RC, Bennett JE, Vogel CL, Carbone PP, DeVita VT. Aspergillosis: the spectrum of the disease in 98 patients. *Medicine (Baltimore)* **1970**;49:147-73.
88. Clarke A, Skelton J, Fraser RS. Fungal tracheobronchitis: report of 9 cases and review of the literature. *Medicine (Baltimore)* **1991**;70:1-14.
89. Just-Nubling G, Ganger G, Hettenkofer M, et al. Therapeutic outcome in AIDS patients with aspergillosis lung infection [abstract PoB09-1409]. In: Program and abstracts of the 9th International Conference on AIDS (Berlin). **1993**.
90. Khoo SH, Bond J, Denning DW. Administering amphotericin B—a practical approach. *J Antimicrob Chemother* **1994**;33:203-13.